

SUTAB INSTRUCTIONS:

The day before your procedure start drinking clear liquids first thing in the morning.

If your procedure is <u>BEFORE</u> 12PM (noon)	If your procedure is <u>AFTER</u> 12PM (noon)
1. At 5:00PM- Open 1 bottle of 12 tablets	1. At 7:00PM- Open 1 bottle of 12 tablets
2. Fill the provided container with 16 ounces of water. Swallow 1 tablet every 2 minutes. You should finish the 12 tablets and entire 16 ounces of water within 25 minutes. *If your experience preparation-related symptoms (nausea, bloating, or cramping), pause or slow the rate of drinking the additional water until symptoms diminish.	2. Fill the provided container with 16 ounces of water. Swallow 1 tablet every 2 minutes. You should finish the 12 tablets and entire 16 ounces of water within 25 minutes. *If your experience preparation-related symptoms (nausea, bloating, or cramping), pause or slow the rate of drinking the additional water until symptoms diminish.
3. Approximately 1 hour after the last tablet is ingested, fill the provided container with 16 ounces of water, and drink the entire amount over 30 minutes.	3. Approximately 1 hour after the last tablet is ingested, fill the provided container with 16 ounces of water, and drink the entire amount over 30 minutes.
4. Approximately 30 minutes after drinking the second container of water, fill the container again with 16 ounces of water and drink the entire amount over 30 minutes.	4. Approximately 30 minutes after drinking the second container of water, fill the container again with 16 ounces of water and drink the entire amount over 30 minutes.
5. At 1:00AM- Open the 2 nd bottle of 12 tablets.	5. At 6:00AM- Open the 2 nd bottle of 12 tablets.
6. Fill the provided container with 16 ounces of water. Swallow 1 tablet every 2 minutes. You should finish the 12 tablets and entire 16 ounces of water within 25 minutes. *If your experience preparation-related symptoms (nausea, bloating, or cramping), pause or slow the rate of drinking the additional water until symptoms diminish.	6. Fill the provided container with 16 ounces of water. Swallow 1 tablet every 2 minutes. You should finish the 12 tablets and entire 16 ounces of water within 25 minutes. *If your experience preparation-related symptoms (nausea, bloating, or cramping), pause or slow the rate of drinking the additional water until symptoms diminish.
7. Approximately 1 hour after the last tablet is ingested, fill the provided container with 16 ounces of water, and drink the entire amount over 30 minutes.	7. Approximately 1 hour after the last tablet is ingested, fill the provided container with 16 ounces of water, and drink the entire amount over 30 minutes.
8. Approximately 30 minutes after drinking the second container of water, fill the container again with 16 ounces of water and drink the entire amount over 30 minutes.	8. Approximately 30 minutes after drinking the second container of water, fill the container again with 16 ounces of water and drink the entire amount over 30 minutes.

DRINK AS MUCH CLEAR LIQUID AS YOU ARE ABLE TO TOLERATE, ALL DAY!
STOP drinking at least 3 hours prior to arrival time.

**** ADMINISTRATIVE FEES**

As a courtesy to other patients who are waiting for appointments, we require that you provide us with notice of any cancellation of an appointment. Missed or cancelled appointments are subject to the following fee, which is not covered by your insurance:

Surgical Center or Hospital Procedure with less than 72 hours' notice.....\$75